



Pneumonia: update on diagnosis and management

Zara Hoare and Wei Shen Lim

BMJ 2006;332:1077-1079
doi:10.1136/bmj.332.7549.1077

Updated information and services can be found at:
<http://bmj.com/cgi/content/full/332/7549/1077>

These include:

References

This article cites 13 articles, 10 of which can be accessed free at:
<http://bmj.com/cgi/content/full/332/7549/1077#BIBL>

1 online articles that cite this article can be accessed at:
<http://bmj.com/cgi/content/full/332/7549/1077#otherarticles>

Rapid responses

5 rapid responses have been posted to this article, which you can access for free at:

<http://bmj.com/cgi/content/full/332/7549/1077#responses>

You can respond to this article at:
<http://bmj.com/cgi/eletter-submit/332/7549/1077>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections

[Other respiratory infections](#) (563 articles)

Notes

To order reprints of this article go to:
<http://www.bmjournals.com/cgi/reprintform>

To subscribe to *BMJ* go to:
<http://bmj.bmjournals.com/subscriptions/subscribe.shtml>

Practice

BMJ Learning

Pneumonia: update on diagnosis and management

Zara Hoare, Wei Shen Lim

Prompt diagnosis and management of community acquired pneumonia saves lives. This article summarises the latest key recommendations in the management of pneumonia and is intended for junior doctors managing this common condition

Why is it important?

Community acquired pneumonia is an important healthcare concern. The annual incidence rate is 6/1000 in the 18-39 age group. This rises to 34/1000 in people aged 75 years and over. Admission to hospital is needed in 20-40% of patients with community acquired pneumonia. About 5-10% of these patients are admitted to intensive care. The overall mortality from community acquired pneumonia is 5-10%, so it is important to identify and treat patients with this disease.^{1,2} We will concentrate on three aspects of hospital management: assessment of disease severity, investigations, and antibiotic treatment.

What are the symptoms and signs of pneumonia?

Pneumonia is defined as an acute lower respiratory tract infection, together with new radiographic shadowing.¹ Anybody can get pneumonia, although it is most common in elderly people. The clinical history of pneumonia may include one or more of:

- Pleuritic chest pain
- Shortness of breath
- Cough
- Production of sputum
- Rigors or night sweats
- Confusion.

On examination the signs may include:

- Raised respiratory rate
- Fever of >38°C
- Focal chest signs: decreased chest expansion, dullness on percussion, decreased entry of air, bronchial breathing, and crackles (none, some, or all of these may be present).

In elderly people, fever may not be present and new onset of mental confusion is more common.³

What investigations should we do?

Table 1 lists the other diagnoses that should be considered when patients present with focal chest signs and increasing shortness of breath. A chest x ray is crucial to making the diagnosis of pneumonia. New chest

Summary points

Pneumonia is common and has high mortality

Simple scoring systems help to distinguish between patients who can be managed at home and those who may need intensive care

Antibiotic treatment is guided by the severity of disease

Newer antibiotics have their place, but penicillin and macrolides (such as erythromycin) are still first line drugs

Editorial by
Goossens and Little



This article is based on a module that is available on BMJ Learning (www.bmjlearning.com)

signs on examination are helpful but not specific. One clinical study conducted in primary care in the United Kingdom showed that only 40% of patients with new lower respiratory tract symptoms and focal chest signs had radiological evidence of pneumonia.⁴

Table 1 Differential diagnosis in community acquired pneumonia

Alternative diagnosis	Supporting clinical feature
Exacerbation of chronic obstructive pulmonary disease	History of the disease or heavy smoking; polyphonic wheeze on examination
Exacerbation of asthma	Patient known to have asthma; wheeze on examination
Pulmonary oedema	Third heart sound and abnormal electrocardiogram
Fibrosing alveolitis	Long history of chronic cough, fine inspiratory crackles on examination
Lung cancer	Long history of chronic cough, haemoptysis, weight loss, and history of smoking
Bronchiectasis	Long history of chronic cough that produces large amounts of purulent sputum
Inhaled foreign body	History of inhalation or coughing when eating
Pulmonary emboli	Sudden onset of shortness of breath and chest pain; occasional haemoptysis

Department of Respiratory Medicine, Nottingham City Hospital, Nottingham NG5 1PB

Zara Hoare
clinical research fellow
of the University of Nottingham

Wei Shen Lim
consultant in general and respiratory medicine

Correspondence to:
Z Hoare
Zara.Hoare@nottingham.ac.uk

BMJ 2006;332:1077-9

About 40% of patients have a pleural effusion, but the amount of fluid is usually small.¹ A microbiological diagnosis can be made on routine microbiological tests in only about 15% of inpatients; this percentage is even lower in patients with mild or moderate pneumonia and no comorbid illnesses.⁵ The following tests are recommended for patients admitted with non-severe community acquired pneumonia:

- Full blood count
- Urea and electrolytes
- Liver function tests
- Sputum culture
- Chest x ray
- Blood cultures.

In patients with severe pneumonia, tests for atypical pathogens (*Mycoplasma pneumoniae*, *Chlamydia* sp, *Coxiella burnetii*, *Legionella* sp) and viruses should also be carried out. These are:

- Legionella urinary antigen test
- Direct immunofluorescence on respiratory samples
- Serological tests on acute and convalescent serum samples.

Testing for pneumococcal urinary antigen is also recommended in severe pneumonia.

How can we identify patients with severe pneumonia?

Severe pneumonia has a high mortality, so it is important to identify patients at high risk of complications and death. Many studies have looked at features associated with poor prognosis. More than 40 indicators of poor prognosis have been identified, and complex scoring systems have been developed.⁶ The most widely studied one is a North American 20 point system,⁷ but this system is not practical in many clinical settings and is not used widely in the UK.


The British Thoracic Society recommends the CURB-65 scoring system.^{8,9} This is a 6 point score (range 0-5) that gives one point each for confusion (abbreviated mental test score ≤ 8 or new disorientation in person, place, or time); urea >7 mmol/l; respiratory rate ≥ 30 /min; low blood pressure (systolic blood pressure <90 mm Hg or diastolic blood pressure ≤ 60 mm Hg); and age 65 years or more. The score relates to mortality, and severe pneumonia is defined as a CURB-65 score of 3 or more. The British Thoracic Society advocates management strategies based on this score (fig 1).

Clinical judgment is needed when using the scoring system. If the score is low but the patient is unwell he or she could still have severe pneumonia. For example, a young person with normal blood pressure and urea but low oxygen saturation despite supplemental oxygen could have severe pneumonia and may need to be reviewed by the intensive care team.

How do we treat it?

Pneumonia must be treated with the appropriate antibiotics. Table 2 shows the current recommendations of the British Thoracic Society for empirical treatment of community acquired pneumonia.

Since the development of new fluoroquinolones with enhanced activity against Gram positive



Assessment of severity of pneumonia with the CURB-65 scoring system. Reproduced and adapted with permission¹

organisms (such as *Streptococcus pneumoniae*), the choice of antibiotic for respiratory tract infections has broadened. Two such agents currently available in the UK are levofloxacin and moxifloxacin. These antimicrobials are active against atypical pathogens and *Streptococcus pneumoniae*. However, concerns exist about the widespread use of these drugs as they have been associated with the emergence of fluoroquinolone resistant streptococci, an increase in the development of methicillin resistant *Staphylococcus aureus* (MRSA), and diarrhoea associated with *Clostridium difficile* infection.^{10,11}

The new fluoroquinolones are not recommended for use in the community but should be reserved for the hospital setting. Currently in the UK, only levofloxacin has a licensed intravenous preparation. This is expensive and not available in many hospitals.

Who should receive intravenous antibiotics?

Patients should receive intravenous antibiotics if they have:

Table 2 Current treatment regimens for pneumonia

Severity of pneumonia	Treatment
Non-severe (home treated or hospital treated) and admitted for non-clinical reasons or previously untreated in the community	Oral amoxicillin 500-1000 mg three times a day, erythromycin 500 mg four times a day, or clarithromycin 500 mg twice a day
Non-severe, hospital treated	Oral amoxicillin 500-1000 mg three times a day plus either erythromycin 500 mg four times a day or clarithromycin 500 mg twice a day. If intravenous treatment is needed, ampicillin 500 mg four times a day or benzylpenicillin 1200 mg four times a day, plus either erythromycin 500 mg four times a day or clarithromycin 500 mg twice a day
Alternative treatment for non-severe hospital treated	Oral levofloxacin 500 mg once daily or moxifloxacin 400 mg once daily
Severe, hospital treated	Intravenous broad spectrum β lactam antibiotics (co-amoxiclav 1.2 g three times a day, cefuroxime 1.5 g three times a day, cefotaxime 1 g three times a day, or ceftriaxone 2 g once daily), plus either intravenous erythromycin 500 mg four times a day or clarithromycin 500 mg twice a day
Alternative treatment for severe, hospital treated	Intravenous levofloxacin* 500 mg twice a day plus benzylpenicillin 1200 mg four times a day

*Other new fluoroquinolones (such as intravenous moxifloxacin) are not currently licensed in severe community acquired pneumonia in the UK.

Sample questions

1. A 70 year old man is admitted to intensive care with severe community acquired pneumonia. Which of the following regimens would you recommend?

- Intravenous cefotaxime 1 g three times a day
- Intravenous cefotaxime 1 g three times a day plus clarithromycin 500 mg twice a day
- Intravenous ceftriaxone 2 g once daily plus erythromycin 250 mg four times a day
- Intravenous co-amoxiclav 1.2 g three times a day plus oral clarithromycin 500 mg twice a day

2. A 52 year old man who smokes and has pneumonia has been taking antibiotics for five days. He is improving clinically. His observations are: temperature 37.3°C, heart rate 80/min, blood pressure 120/80 mm Hg, respiratory rate 14/min. Which of these statements are correct?

- It is appropriate to discharge him home
- He should stay in hospital
- He should stop his antibiotics
- He does not need a repeat chest x ray
- He needs to be seen in clinic in two weeks

- Severe pneumonia (CURB-65 score ≥ 3)
- Functional or anatomical reasons for malabsorption
- Impaired consciousness
- Impaired swallowing reflex.¹

Intravenous antibiotics should be changed to oral as soon as possible. Resolution of the patient's fever, improved oxygen status, and haemodynamic stability are indicators for oral treatment.

In non-severe pneumonia, antibiotics should be given for seven days. Patients with severe pneumonia but no microbiological confirmation usually receive 10 days of intravenous drugs. Patients who are infected with *Legionella* sp, *S aureus*, and Gram negative bacteria should receive 14-21 days of antimicrobial treatment.

C reactive protein

Measuring C reactive protein (an acute phase protein) concentrations is useful for managing pneumonia. Ninety five percent of patients admitted with community acquired pneumonia have a C reactive protein value > 50 mg/l.¹² The initial concentration of this protein does not correlate closely with the severity of disease, but it can be used as an indicator of response to treatment. Treatment failure or complications (such as an empyema) should be considered if C reactive protein is not 50% lower by the fourth day.¹³

When should patients be discharged home?

It is important to know when it is safe to discharge patients. Features of clinical instability are:

- Temperature $> 37.8^\circ\text{C}$
- Heart rate $> 100/\text{min}$
- Respiratory rate $> 24/\text{min}$
- Systolic blood pressure < 90 mm Hg
- Oxygen saturation $< 90\%$

- Abnormal mental status
- Inability to take oral drugs.

One study showed that 40% of patients who had two or more of these features within 24 hours before they were discharged died or were readmitted within 30 days, compared with 11% of those with none of these features.¹⁴

Every patient with pneumonia should be reviewed about six weeks after discharge. In patients with poor clinical recovery, a chest x ray should be done to identify those with possible underlying malignancy. Patients with a high risk of malignancy, such as those who smoke or who are aged over 50 years, should also have a repeat chest x ray at six weeks.

Competing interests: None declared.

- 1 British Thoracic Society. Guidelines for the management of community acquired pneumonia in adults. *Thorax* 2001;56(suppl 4):1-64.
- 2 Macfarlane JT, Boldy D. 2004 update of BTS guidelines: what's new? *Thorax* 2004;59:364-6.
- 3 Metlay JP, Schulz R, Li YH, Singer DE, Marrie TJ, Coley CM, et al. Influence of age on symptoms at presentation in patients with community-acquired pneumonia in the community. *Arch Intern Med* 1997;157:1453-9.
- 4 Woodhead MA, Macfarlane JT, McCracken JS, Rose DH, Finch RG. Prospective study of the aetiology and outcome of pneumonia in the community. *Lancet* 1987;1:671-4.
- 5 Woodhead MA, Arrowsmith J, Chamberlain-Webber R, Wooding S, Williams I. The value of routine microbial investigation in community-acquired pneumonia. *Respir Med* 1991;85:313-7.
- 6 Metlay JP, Fine MJ. Testing strategies in the initial management of patients with community-acquired pneumonia. *Ann Intern Med* 2003;138:109-18.
- 7 Fine MJ, Auble TE, Yearly DM, Hanuse BH, Weissfeld LA, Singer DE, et al. A prediction rule to identify low-risk patients with community-acquired pneumonia. *N Engl J Med* 1997;336:243-50.
- 8 Lim WS, Van der Eerden MM, Laing R, Boersma WG, Karalus N, Town GI, et al. Defining community acquired pneumonia severity on presentation to hospital: an international derivation and validation study. *Thorax* 2003;58:377-82.
- 9 Ewig S, Torres A, Woodhead M. Assessment of pneumonia severity: a European perspective. *Eur Respir J* 2006;27:6-8.
- 10 Weber SG, Gold HS, Hoopers DC, Karchmer AW, Carmeoli Y. Fluoroquinolones and the risk of methicillin-resistant *Staphylococcus aureus* in hospitalized patients. *Emerg Infect Dis* 2003;9:1415-22.
- 11 Loo VE, Poirer L, Miller MA, Oughton M, Libman MD, Michaud S, et al. A predominantly clonal multi-institutional outbreak of *Clostridium difficile*-associated diarrhea with high morbidity and mortality. *N Engl J Med* 2005;353:2442-9.
- 12 Hansson LO, Hedlund JU, Ortqvist A. Sequential changes of inflammatory and nutritional markers in patients with community acquired pneumonia. *Scand J Clin Lab Invest* 1997;57:11-8.
- 13 Smith RP, Lipworth BJ, Cree IA, Spiers EM, Winter JH, et al. C-reactive protein. A clinical marker in community acquired pneumonia. *Chest* 1995;108:1288-91.
- 14 Halm EA, Fine MJ, Kapoor WN, Singer DE, Marrie TJ, Siu AC. Instability on hospital discharge and the risk of adverse outcome in patients with pneumonia. *Arch Intern Med* 2002;162:1278-84.

Interactive case report**70 year old man with scrotal swelling, abdominal aortic aneurysm, and renal lesions**

This case was described on 15 and 22 April (*BMJ* 2006;332:899, 959). Debate on the patient's management continues on bmj.com (<http://bmj.bmjournals.com/cgi/eletters/332/7546/899>). On 13 May we will publish the case outcome together with commentaries on the issues raised by the management and online discussion from relevant experts and the patient.